

# Lesson 1

Background:

## Herbicide Resistance in Aquatics and Description of Aquatic Plant Management Venues and Plant Types



# Objectives:

By the end of this lesson you will:



Recognize key differences between aquatic plant control and crop management that effect herbicide use in water

# Objectives:

By the end of this lesson you will:



Recognize key differences between aquatic plant control and crop management that effect herbicide use in water



Understand how the uses or functions of different types of aquatic systems influence control strategies

# Objectives:

By the end of this lesson you will:



Recognize key differences between aquatic plant control and crop management that effect herbicide use in water



Understand how the uses or functions of different types of aquatic systems influence control strategies



Learn how plant types influence selection of herbicide mode, amount, and application frequency

# Uses and Functions of Aquatic Systems:

The uses and functions of aquatic systems greatly influence plant management strategies within these waters. Waters can generally be divided into **Man-made** systems and **Natural** or modified natural systems.

## Man-made systems:

- include water storage basins, conveyance canals for flood control, irrigation, and potable water supply, water retention ponds



## Man-made systems:

- include water storage basins, conveyance canals for flood control, irrigation, and potable water supply, water retention ponds
- any amount of aquatic macrophyte growth may be considered as undesirable



## Man-made systems:

- include water storage basins, conveyance canals for flood control, irrigation, and potable water supply, water retention ponds
- any amount of aquatic macrophyte growth may be considered as undesirable
- are similar to commercial crop herbicide application strategies
  - plants are generally subject to maximum control efforts with low emphasis on selectivity



## Man-made systems:

- include water storage basins, conveyance control, irrigation, and potable
- any amount of

Note: Man-made systems also include thousands of water feature ponds in which non-desirable plants and algae are intensively managed. While technically man-made, these systems are managed for aesthetic reasons and often to promote non-target plants and animals.



## Natural systems:

- include ponds, lakes, rivers, and reservoirs



## Natural systems:

- include ponds, lakes, rivers, and reservoirs
- with the exception of invasive plants, most plant and animal species may be considered valuable ecosystem components



## Natural systems:

- include ponds, lakes, rivers, and reservoirs
- with the exception of invasive plants, most plant and animal species may be considered valuable ecosystem components
- control strategies need to effectively control 1-2 target plants while conserving multiple native or other valuable species



## Natural systems:

- include ponds, lakes, rivers, and reservoirs
- with the exception of invasive plants, most plant and animal species may be considered valuable ecosystem components
- control strategies need to effectively control 1-2 target plants while conserving multiple native or other valuable species
- selectivity considerations may:
  - limit available herbicide active ingredients
  - limit rates and timing of herbicide applications
  - influence cost and amount of control effort



## Key differences between crop management and aquatic plant control, especially in natural systems:

In crop management, many weeds are targeted for control, usually among one or two non-target crop species

## Key differences between crop management and aquatic plant control, especially in natural systems:

In crop management, many weeds are targeted for control, usually among one or two non-target crop species

In natural areas, one or two plants, usually invasive plants, are targeted among many desirable plant and animal species

## Key differences between crop management and aquatic plant control, especially in natural systems:

In crop management, many weeds are targeted for control, usually among one or two non-target crop species

In natural areas, one or two plants, usually invasive plants, are targeted among many desirable plant and animal species



Conserving or enhancing desirable species is equally or more important than control of a target plant

# Plant Types in Aquatic Systems Include:

Emergent  
Floating  
Submersed  
Algae

# Emergent Plants



- include grasses, sedges and rushes and numerous broadleaf species

# Emergent Plants



- include grasses, sedges and rushes and numerous broadleaf species
- control, especially large-scale control, is similar to commercial crop management

# Emergent Plants



- include grasses, sedges and rushes and numerous broadleaf species
- control, especially large-scale control, is similar to commercial crop management
  - **vast area monocultures may be targeted**
  - herbicides are applied directly to plant foliage
  - herbicides can be precisely applied to a defined area

# Emergent Plants



- include grasses, sedges and rushes and numerous broadleaf species
- control, especially large-scale control, is similar to commercial crop management
  - vast area monocultures may be targeted
  - **herbicides are applied directly to plant foliage**
  - herbicides can be precisely applied to a defined area

# Emergent Plants



- include grasses, sedges and rushes and numerous broadleaf species
- control, especially large-scale control, is similar to commercial crop management
  - vast area monocultures may be targeted
  - herbicides are applied directly to plant foliage
  - **herbicides can be precisely applied to a defined area**

# Emergent Plants



- include grasses, sedges and rushes and numerous broadleaf species
- control, especially large-scale control, is similar to commercial crop management
  - vast area monocultures may be targeted
  - herbicides are applied directly to plant foliage
  - herbicides can be precisely applied to a defined area
- selectivity concerns may reduce the number of available herbicides
  - i.e., when controlling invasive grasses growing among native grasses or other desirable species

# Floating Plants



- include large water hyacinth and water lettuce and minute species like salvinia, duckweeds, and watermeal

# Floating Plants



- include large water hyacinth and water lettuce and minute species like salvinia, duckweeds, and watermeal
- herbicides often applied directly to foliage, similar to emergent plant and commercial crop management
  - frequent spot applications for selective control among more desirable plant species

# Floating Plants



- include large water hyacinth and water lettuce and minute species like salvinia, duckweeds, and watermeal
- herbicides often applied directly to foliage, similar to emergent plant and commercial crop management
  - frequent spot applications for selective control among more desirable plant species
- injecting herbicides into water for root uptake is becoming more common

# Submersed Plants



- include invasive plants like hydrilla and Eurasian watermilfoil in natural systems

# Submersed Plants




- include invasive plants like hydrilla and Eurasian watermilfoil in natural systems
- pondweeds and other species in water conveyance systems

# Submersed Plants



- include invasive plants like hydrilla and Eurasian watermilfoil in natural systems
- pondweeds and other species in water conveyance systems
- native submersed plants may be managed in natural areas, especially in late season for access and recreation



Late season native Illinois pondweed growth in 12,000-acre East Lake Toho, FL



# Submersed Plant Herbicide Applications

Most strikingly different from commercial crop management

# Submersed Plant Herbicide Applications

**Most strikingly different from commercial crop management**

- entire water column is usually treated in submersed plant control vs. foliar applications in row crop maintenance

# Submersed Plant Herbicide Applications

## Most strikingly different from commercial crop management

- entire water column is usually treated in submersed plant control vs. foliar applications in row crop maintenance
- herbicide exposure often measured in hours in crop management

# Submersed Plant Herbicide Applications

## Most strikingly different from commercial crop management

- entire water column is usually treated in submersed plant control vs. foliar applications in row crop maintenance
- herbicide exposure often measured in hours in crop management
  - hours / days for aquatic contact herbicides, and

# Submersed Plant Herbicide Applications

## Most strikingly different from commercial crop management

- entire water column is usually treated in submersed plant control vs. foliar applications in row crop maintenance
- herbicide exposure often measured in hours in crop management
  - hours / days for aquatic contact herbicides, and
  - weeks / months for aquatic systemic herbicides

# Submersed Plant Herbicide Applications

## Most strikingly different from commercial crop management

- entire water column is usually treated in submersed plant control vs. foliar applications in row crop maintenance
- herbicide exposure often measured in hours in crop management
  - hours / days for aquatic contact herbicides, and
  - weeks / months for aquatic systemic herbicides
- applied in systems where water is constantly moving

# Submersed Plant Herbicide Applications

## Most strikingly different from commercial crop management

- entire water column is usually treated in submersed plant control vs. foliar applications in row crop maintenance
- herbicide exposure often measured in hours in crop management
  - hours / days for aquatic contact herbicides, and
  - weeks / months for aquatic systemic herbicides
- applied in systems where water is constantly moving
- herbicides dissipate immediately upon application to water

# Submersed Plant Herbicide Applications

## Most strikingly different from commercial crop management


- entire water column is usually treated in submersed plant control vs. foliar applications in row crop maintenance
- herbicide exposure often measured in hours in crop management
  - hours / days for aquatic contact herbicides, and
  - weeks / months for aquatic systemic herbicides
- applied in systems where water is constantly moving
- herbicides dissipate immediately upon application to water
- management objectives usually include controlling invasive species while enhancing many comingled non-target species

# Submersed Plant Herbicide Applications

## Most strikingly different from commercial crop management


- entire water column is usually treated in submersed plant control vs. foliar applications in row crop maintenance
- herbicide exposure often measured in hours in crop management
  - hours / days for aquatic contact herbicides, and
  - weeks / months for aquatic systemic herbicides
- applied in systems where water is constantly moving
- herbicides dissipate immediately upon application to water
- management objectives usually include controlling invasive species while enhancing many comingled non-target species
- pre-emergent strategies are rarely applicable for submersed plant control

# Algae



- include filamentous, planktonic, blue-green, and macrophytic algae

# Algae



- include filamentous, planktonic, blue-green, and macrophytic algae
- blue-green algae (cyanobacteria) can produce toxins, cause taste and odor problems in drinking water

# Algae



- include filamentous, planktonic, blue-green, and macrophytic algae
- blue-green algae (cyanobacteria) can produce toxins, cause taste and odor problems in drinking water
- copper-based compounds have been the dominant mode of action to control algae for decades

# Algae



- include filamentous, planktonic, blue-green, and macrophytic algae
- blue-green algae (cyanobacteria) can produce toxins, cause taste and odor problems in drinking water
- copper-based compounds have been the dominant mode of action to control algae for decades
- few realistic alternatives to copper

# Algae



- include filamentous, planktonic, blue-green, and macrophytic algae
- blue-green algae (cyanobacteria) can produce toxins, cause taste and odor problems in drinking water
- copper-based compounds have been the dominant mode of action to control algae for decades
- few realistic alternatives to copper
  - most aquatic herbicides do not have algal activity at label rates

# Key Considerations from Lesson 1:

- In crop management, many weeds are targeted for control, usually among one or two non-target crop species

# Key Considerations from Lesson 1:

- In crop management, many weeds are targeted for control, usually among one or two non-target crop species
- In natural aquatic systems, one or two plants, usually invasive plants, are targeted among many desirable plants and animals

# Key Considerations from Lesson 1:

- In crop management, many weeds are targeted for control, usually among one or two non-target crop species
- In natural aquatic systems, one or two plants, usually invasive plants, are targeted among many desirable plants and animals
- Submersed plant control is most strikingly different from commercial crop management

# Key Considerations from Lesson 1:

- In crop management, many weeds are targeted for control, usually among one or two non-target crop species
- In natural aquatic systems, one or two plants, usually invasive plants, are targeted among many desirable plants and animals
- Submersed plant control is most strikingly different from commercial crop management
- Conserving or enhancing desirable species is equally or more important than control of a target plant in natural aquatic systems